

AS Pure Pre-mock Nov 2022

This exam has 15 questions, for a total of 100 marks.

- Print in “booklets” will allow all questions to be on the left hand side.

Question	Marks	Score
1	5	
2	6	
3	6	
4	7	
5	6	
6	6	
7	5	
8	5	
9	6	
10	7	
11	9	
12	10	
13	7	
14	4	
15	11	
Total:	100	

2. [In this question the unit vectors \mathbf{i} and \mathbf{j} are due east and due north respectively]

A boat B is moving with constant velocity. At noon, B is at the point with position vector $(3\mathbf{i} - 4\mathbf{j})$ km with respect to a fixed origin O . At 1430 on the same day, B is at the point with position vector $(8\mathbf{i} + 11\mathbf{j})$ km.

(a) Calculate the bearing on which the boat is moving. (3)

(b) Calculate the speed of the boat, giving your answer in km h^{-1} (3)

**In this question you should show all stages of your working.
Solutions relying on calculator technology are not acceptable.**

3. (i) Solve the equation

$$4^{2x+1} = 8^{4x}$$

(3)

(ii) Solve

$$3\sqrt{18} - \sqrt{32} = \sqrt{n}$$

(3)

4. A bakery makes pies.

On any day, the total cost to the bakery, $\pounds y$, of making x pies is modelled to be the sum of two separate elements:

- a fixed cost, C ,
- a cost that is proportional to the number of pies that are made that day, K .

(a) Write down a general equation linking y with x , for this model. (1)

The pies are sold for $\pounds 5$ each.

On a day when 650 pies are made and sold, the bakery makes a profit of $\pounds 200$.

On a day when 230 pies are made and sold, the bakery makes a loss of $\pounds 80$.

Using the above information,


(b) Rewrite your answer to part (a), giving the values of the constants C and K as simplified fractions. (3)

(c) With reference to the model, interpret the significance of the value for the gradient in the equation derived above. (1)

Assuming that each pie is sold on the day it is made,

(d) find the least number of pies that must be made on any given day for the bakery to make a profit that day. (2)

5.



Not to scale

Figure 1

Figure 1 shows the design for a structure used to support a roof.

The structure consists of four wooden beams, AB , BD , BC , AD .

- $AB = 6.5$ m
- $BC = BD = 4.7$ m
- $\angle BAC = 35^\circ$

(a) Find, to one decimal place, the size of $\angle ACB$.

(3)

(b) Find, to the nearest metre, the total length of wood required to make this structure.

(3)

6. (a) Find the first 3 terms in ascending powers of x of the binomial expansion of

$$(2 + ax)^6$$

where a is a non-zero constant. Give each term in simplest form.

(4)

Given that, in the expansion, the coefficient of x is equal to the coefficient of x^2 .

(b) Find the value of a .

(2)

Question 6 continued

Lined writing area for the answer to Question 6.

(Total for Question 6 is 6 marks)

9. A population of a rare species of toad is being studied.

The number of toads, N , in the population, t years after the start of the study, is modelled by the equation.

$$N = \frac{900e^{0.12t}}{2e^{0.12t} + 1} \quad t \geq 0, t \in \mathbb{R}$$

According to this model,

- (a) Calculate the number of toads in the population at the start of the study. (1)
- (b) Find the value of t when there are 420 toads in the population, giving your answer to 2 decimal places. (4)
- (c) Explain why, according to this model, the number of toads can never reach 500. (1)

10.




Figure 2

Figure 2 shows a plot of the curve with equation $y = \sin \theta$, $0 \leq \theta \leq 360^\circ$

- (a) State the coordinates of the minimum point on the curve with equation

$$y = 4 \sin \theta, \quad 0 \leq \theta \leq 360^\circ$$

(2)

A copy of Figure 2, called Diagram 1, is shown on the next page.

- (b) On Diagram 1, sketch and label the curves

(i) $y = 1 + \sin \theta$, $0 \leq \theta \leq 360^\circ$

(ii) $y = \tan \theta$, $0 \leq \theta \leq 360^\circ$

(2)

- (c) Hence find the number of solutions of the equation

(i) $\tan \theta = 1 + \sin \theta$, $0 \leq \theta \leq 2160^\circ$

(ii) $\tan \theta = 1 + \sin \theta$, $0 \leq \theta \leq 1980^\circ$

(3)

Question 10 continued




Diagram 1

(Total for Question 10 is 7 marks)

12. (i) A circle C_1 has centre $A(2, 1)$ and passes through the point $B(10, 7)$.

The line l is the tangent to C_1 at the point B .

Find an equation of l in the form $ax + by + c = 0$, where a , b and c are integers to be found.

(5)

- (ii) A different circle C_2 has equation

$$x^2 + y^2 - 6x - 14y + k = 0$$

where k is a constant.

Given that C_2 lies entirely in the first quadrant, find the range of possible values for k .

(5)

13. A doctors' surgery starts a campaign to reduce missed appointments. The number of missed appointments for each of the first five weeks after the start of the campaign is shown below.

Number of weeks after the start (x)	1	2	3	4	5
Number of missed appointments (y)	235	149	99	59	38

Figure 3

This data could be modelled by an equation of the form $y = pq^x$ where p and q are constants.

- (a) Show that this relationship may be expressed in the form $\log_{10} y = mx + c$, expressing m and c in terms of p and/or q . (2)




Figure 4

Figure 4 shows $\log_{10} y$ plotted against x , for the given data.

- (b) Estimate the values of p and q . (3)
- (c) Use the model to predict when the number of missed appointments will fall below 20. Explain why this answer may not be reliable. (2)

14. Prove that for all $n \in \mathbb{N}$, $n^2 - 2n + 2$ is not divisible by 4.

(4)

SohokMaths by A. Chan sohokmaths.com

15. A curve has equation $y = f(x)$, where

$$f''(x) = \frac{6}{\sqrt{x^3}} + x \quad x > 0$$

The point $P(4, -50)$ lies on the curve.

Given that $f'(x) = -4$ at P ,

- (a) Find the equation of the normal at P , write your answer in the form $y = mx + c$, where m and c are constants.

(3)

- (b) Find $f(x)$

(8)
